



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12N 15/00, 1/21, 15/11 A61K 39/00	A1	(11) International Publication Number: WO 92/01783 (43) International Publication Date: 6 February 1992 (06.02.92)
(21) International Application Number: PCT/US91/04833 (22) International Filing Date: 9 July 1991 (09.07.91) (30) Priority data: 553,907 16 July 1990 (16.07.90) US (71) Applicants: ALBERT EINSTEIN COLLEGE OF MEDICINE OF YESHIVA UNIVERSITY [US/US]; 1300 Morris Park Avenue, Bronx, NY 10461 (US). UNIVERSITY OF PITTSBURGH [US/US]; 911 William Pitt Union, Pittsburgh, PA 15260 (US). (72) Inventors: JACOBS, William, R. ; 163 Fordham Street, City Island, NY 10464 (US). HATFULL, Graham ; 5 Forbes Terrace, Pittsburgh, PA 15217 (US).		(74) Agents: OLSTEIN, Elliot, M. et al.; Carella, Byrne, Bain, Gilfillan, Cecchi & Stewart, 6 Becker Farm Road, Roseland, NJ 07068 (US). (81) Designated States: AT (European patent), AU, BE (European patent), BR, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent). Published <i>With international search report.</i>
(54) Title: DNA CAPABLE OF SITE-SPECIFIC INTEGRATION INTO MYCOBACTERIA (57) Abstract A DNA for integrating DNA into a mycobacterium chromosome portion which includes a first DNA sequence which is a phage DNA portion encoding bacteriophage integration into a mycobacterium chromosome, and a second DNA sequence encoding a protein or polypeptide heterologous to the mycobacterium in which the DNA is to be integrated. Such DNA may be integrated into mycobacteria, which may then be administered as a vaccine and/or therapeutic agent.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CC	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU ⁺	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark				

⁺ It is not yet known for which States of the former Soviet Union any designation of the Soviet Union has effect.

-1-

**DNA CAPABLE OF SITE-SPECIFIC INTEGRATION
INTO MYCOBACTERIA**

This invention relates to DNA capable of integrating into mycobacterial chromosomes. More particularly, this invention relates to DNA which is capable of site-specific integration into a mycobacterium chromosome while containing a DNA sequence encoding a protein heterologous to the mycobacterium in which the DNA is integrated.

Certain mycobacteria represent major pathogens of man and animals. For example, tuberculosis is generally caused in humans by Mycobacterium tuberculosis, and in cattle by Mycobacterium bovis, which may also be transmitted to humans and other animals. Mycobacteria leprae is the causative agent of leprosy. M. tuberculosis and mycobacteria of the avium-intracellulare-scrofulaceum group (MAIS group) represent major opportunistic pathogens of patients with acquired immune deficiency syndrome (AIDS). M. pseudotuberculosis is a major pathogen of cattle.

On the other hand, Bacille Calmette-Guerin, or BCG, an avirulent strain of M. bovis, is widely used in human vaccines, and in particular is used as a live vaccine, which is protective against tuberculosis. BCG is the only childhood vaccine which is currently given at birth, has a very low incidence of adverse effects, and can be used repeatedly in an individual (eg., in multiple forms). In addition, BCG and other mycobacteria (eg.,

M. smegmatis), employed in vaccines, have adjuvant properties among the best currently known and, therefore, stimulate a recipient's immune system to respond to antigens with great effectiveness.

It has been suggested by Jacobs, et al, Nature, Vol. 327, No. 6122, pgs. 532-535 (June 11, 1987) that BCG could be used as a host for the construction of recombinant vaccines. In other words, it was suggested to take an existing vaccine (in this case against tuberculosis) and expand its protective repertoire through the introduction of one or more genes from other pathogens. Because BCG vaccines are administered as live bacteria, it is essential that any foreign antigens, polypeptides, or proteins expressed by the bacteria are not lost from the bacteria subsequent to vaccination.

Transformation, the process whereby naked DNA is introduced into bacterial cells, has been carried out successfully in mycobacteria. Jacobs, et al (1987), hereinabove cited, have described transformation of mycobacteria through chemical methods, and Snapper, et al. PNAS, Vol. 85, pgs. 6987-6991 (September 1988) have described transformation of mycobacteria by electroporation. Electroporation can give from 10^5 to 10^6 transformants per μg of plasmid DNA and such plasmid DNA's may carry genes for resistance to antibiotic markers such as kanamycin (Snapper, et al 1988) to allow for selection of transformed cells from non-transformed cells.

Jacobs, et al (1987) and Snapper, et al (1988) have also described the use of cloning vehicles, such as plasmids and bacteriophages, for carrying genes of interest into mycobacteria.

Combination of the above-mentioned techniques, along with standard tools of molecular cloning (e.g., use of restriction enzymes, etc.) allows the cloning of genes of interest into vectors and introduction of such genes into mycobacteria. To express these genes, it is important to have available signals for gene expression, in particular, transcription promoter

elements. Such promoter elements have been isolated from mycobacterial heat shock genes, and used to express foreign antigens in mycobacteria.

Plasmids currently available for use in mycobacteria, however, are not stably maintained, and are readily lost during non-selective growth. If recombinant mycobacteria expressing genes of interest are to be employed as vaccines against a particular antigen or as a means of providing a therapeutic agent to a host, which is expressed by the mycobacteria, it is crucial that such genes not be lost from the recombinant mycobacteria subsequent to their administration.

It is an object of the present invention to introduce into mycobacteria and ultimately express a variety of heterologous genes, including, but not limited to, genes for protective antigen(s) for a variety of pathogens, and/or for other therapeutic agents, thereby enabling one to produce effective vaccines against such pathogens and/or to provide effective therapeutic agents, whereby such genes will not be lost subsequent to vaccination or administration of the therapeutic agent.

In accordance with an aspect of the present invention, there is provided a DNA which comprises a first DNA sequence which is a phage DNA portion encoding bacteriophage integration into a mycobacterium chromosome, and a second DNA sequence encoding at least one protein or polypeptide which is heterologous to the mycobacterium in which the DNA is to be integrated.

The term "phage DNA portion", as used herein, means that the DNA sequence is derived from a phage and lacks the DNA which is required for phage replication.

Bacteriophages from which the phage DNA portion may be derived include, but are not limited to, mycobacteriophages, such as but not limited to the L5, L1, Bxb1 and TM4 mycobacteriophages; the lambda phage of E.coli; the toxin phages

of Corynebacteria; phages of Actinomycetes and Norcadia, the ϕ C31 phage of Streptomyces; and the P22 phage of Salmonella. Preferably, the phage DNA portion encodes mycobacteriophage integration into a mycobacterium chromosome.

In a preferred embodiment, the first DNA sequence includes DNA encoding integrase, which is a protein that provides for integration of the DNA into the mycobacterial chromosome. Most preferably, the first DNA sequence also includes DNA which encodes an AttP site.

The DNA sequence encoding the AttP site and the integrase provides for an integration event which is referred to as site-specific integration. DNA containing the AttP site and the integrase gene is capable of integration into a corresponding AttB site of a mycobacterium chromosome.

It is to be understood that the exact DNA sequence encoding the attP site may vary among different phages, and that the exact DNA sequence encoding the attB site may vary among different mycobacteria.

The integration event results in the formation of two new junction sites called AttL and AttR, each of which contain part of each of AttP and AttB. The inserted and integrated non-phage DNA which includes the first and second DNA sequences, is flanked by the AttL and AttR sites. The insertion and integration of the phage DNA portion results in the formation of a transformed mycobacterium.

Applicants have found that, when employing the phage DNA portion of the present invention for integration into mycobacterial chromosome, the gene(s) of interest which is integrated into the mycobacterial chromosome is not lost following non-selective growth of the mycobacteria. Thus, the gene(s) of interest may be expressed by the mycobacteria following such non-selective growth, thus making such transformed mycobacteria excellent vehicles to be employed in vaccines or pharmaceuticals whereby such mycobacteria will express antigens

and/or therapeutic agents of interest subsequent to administration of the recombinant mycobacteria to a host.

Mycobacteria into which the phage DNA portion may be integrated include, but are not limited to, Mycobacterium bovis-BCG, M. smegmatis, M. avium, M. phlei, M. fortuitum, M. lufu, M. paratuberculosis, M. habana, M. scrofalaceum, M. leprae and M. intracellulare. In a preferred embodiment, the DNA is integrated into Mycobacterium bovis- BCG.

The second DNA sequence which encodes a protein heterologous to mycobacteria may be DNA which is all or a portion of a gene encoding protein(s) or polypeptide(s) of interest; DNA encoding a selectable marker or markers; or DNA encoding both a selectable marker or markers and at least one protein or polypeptide of interest.

Proteins or polypeptides of interest, which may be encoded by the second DNA sequence include, but are not limited to, antigens, anti-tumor agents, enzymes, lymphokines, pharmacologic agents, immunopotentiators, and reporter molecules of interest in a diagnostic context.

Antigens for which the second DNA sequence may encode include, but are not limited to, Mycobacterium leprae antigens; Mycobacterium tuberculosis antigens; Rickettsia antigens; malaria sporozoites and merozoites; diphtheria toxoids; tetanus toxoids; Clostridium antigens; Leishmania antigens; Salmonella antigens; Borrelia antigens; Mycobacterium africanum antigens; Mycobacterium intracellulare antigens; Mycobacterium avium antigens; Treponema antigens; Pertussis antigens; Schistosoma antigens; Filaria antigens; Herpes virus antigens; influenza and parainfluenza virus antigens; measles virus antigens; mumps virus antigens; hepatitis virus antigens; Shigella antigens; Neisseria antigens; rabies antigens, polio virus antigens; Rift Valley Fever virus antigens; dengue virus antigens; measles virus antigens; Human Immunodeficiency Virus (HIV) antigens; respiratory syncytial virus (RSV) antigens; snake venom antigens;

and Vibrio cholera antigens. Enzymes which may be encoded include, but are not limited to, steroid enzymes.

Anti-tumor agents which may be encoded by the second DNA sequence include, but are not limited to, interferon- α , interferon- β , or interferon- γ , and tumor necrosis factor, or TNF. Lymphokines which may be encoded include, but are not limited to, interleukins 1 through 8.

Reporter molecules which may be encoded include, but are not limited to, luciferase, B-galactosidase, B-glucuronidase, and catechol dehydrogenase.

Other peptides or proteins which may be encoded by the second DNA sequence include, but are not limited to, those which encode for stress proteins, which can be administered to evoke an immune response or to induce tolerance in an autoimmune disease (e.g., rheumatoid arthritis).

Selectable markers which may be encoded include, but are not limited to, the kanamycin resistance marker, the neomycin resistance marker, the chloroamphenicol resistance marker, and the hygromycin resistance marker.

The phage DNA portion of the present invention, which includes the first DNA sequence encoding mycobacterium phage integration into a mycobacterium chromosome, and the second DNA sequence encoding at least one protein or polypeptide heterologous to mycobacteria, may be constructed through genetic engineering techniques known to those skilled in the art. In a preferred embodiment, the phage DNA portion may be a plasmid including, in addition to the DNA encoding integration and the DNA encoding a heterologous protein, an origin of replication for any of a wide variety of organisms, which includes, but is not limited to, E.coli, Streptomyces species, Bacillus species, Staphylococcus species, Shigella species, Salmonella species and various species of pneumococci. Most preferably, the plasmid includes an origin of replication for E.coli.

The phage DNA portion also may include a suitable promoter. Suitable promoters include, but are not limited to, mycobacterial promoters such as the BCG HSP60 and HSP70 promoters; mycobactin promoters of M. tuberculosis and BCG, the superoxide dismutase promoter, the α -antigen promoter of M. tuberculosis and BCG, the MBP-70 promoter, the 45 kda antigen promoter of M. tuberculosis and BCG; and the mycobacterial asd promoter; the mycobacterial 14 kda and 12 kda antigen promoters; mycobacteriophage promoters such as the Bxb1 promoter, the L1 and L5 promoters, and the TM4 promoters; E.coli promoters; or any other suitable promoter. The selection of a suitable promoter is deemed to be within the scope of those of ordinary skill in the art from the teachings contained herein.

The promoter sequence may, in one embodiment, be part of an expression cassette which also includes a portion of the gene normally under the control of the promoter. For example, when a mycobacterial HSP60 or HSP70 promoter is employed, the expression cassette may include, in addition to the promoter, a portion of the gene for the HSP60 or HSP70 protein. When the expression cassette and the DNA encoding the heterologous protein or polypeptide are expressed, the protein expressed by the cassette and the DNA encoding the heterologous protein or polypeptide is a fusion protein of a fragment of a mycobacterial protein (eg., the HSP60 or HSP70 protein), and the heterologous protein.

In a preferred embodiment, the transcription initiation site, the ribosomal binding site, and the start codon, which provides for the initiation of the translation of mRNA, are each of mycobacterial origin. The stop codon, which stops translation of mRNA, thereby terminating synthesis of the heterologous protein, and the transcription termination site, may be of mycobacterial origin, or of other bacterial origin, or such stop codon and transcription termination site may be those of the DNA encoding the heterologous protein or polypeptide.

The phage DNA portion may be employed, as hereinabove indicated, for integration into a mycobacterial chromosome, thereby transforming the mycobacteria, and whereby the mycobacteria will express protein(s) or polypeptide(s) heterologous to mycobacteria. Such mycobacteria may be utilized in the production of a vaccine or a therapeutic agent, depending upon the protein(s) or polypeptides expressed by the transformed mycobacteria. To form such a vaccine or therapeutic agent, the transformed mycobacteria are administered in conjunction with a suitable pharmaceutical carrier. As representative examples of suitable carriers there may be mentioned: mineral oil, alum, synthetic polymers, etc. Vehicles for vaccines and therapeutic agents are well known in the art and the selection of a suitable vehicle is deemed to be within the scope of those skilled in the art from the teachings contained herein. The selection of a suitable vehicle is also dependent upon the manner in which the vaccine or therapeutic agent is to be administered. The vaccine or therapeutic agent may be in the form of an injectable dose and may be administered intramuscularly, intravenously, orally, intradermally, or by subcutaneous administration.

Other means for administering the vaccine or therapeutic agent should be apparent to those skilled in the art from the teachings herein; accordingly, the scope of the invention is not to be limited to a particular delivery form.

When the transformed mycobacteria are employed as a vaccine, such a vaccine has important advantages over other presently available vaccines. Mycobacteria have, as hereinabove indicated, adjuvant properties among the best currently known and, therefore, stimulate a recipient's immune system to respond with great effectiveness. This aspect of the vaccine induces cell-mediated immunity and thus is especially useful in providing immunity against pathogens in cases where cell-mediated immunity appears to be critical for resistance. Also, mycobacteria may stimulate long-term memory or immunity. It thus may be possible

to prime long-lasting T cell memory, which stimulates secondary antibody responses neutralizing to the infectious agent or the toxin. Such priming of T cell memory is useful, for example, against tetanus and diphtheria toxins, pertussis, malaria, influenza virus, Herpes virus, rabies, Rift Valley fever virus, dengue virus, measles virus, Human Immunodeficiency Virus (HIV), respiratory syncytial virus, human tumors, and snake venoms. Another advantage in employing mycobacteria transformed with the phage DNA portion of the present invention as a vaccine or a therapeutic agent is that mycobacteria in general have a large genome (i.e., approximately 3×10^6 base pairs in length). Because the genome is large, it is able to accommodate a large amount of DNA from other source(s), and may possibly be employed to make a vaccine and/or therapeutic agent containing DNA sequences encoding more than one antigen and/or therapeutic agent.

Also, because the integrated gene(s) of interest has not been lost following non-selective growth of transformed mycobacteria, the gene(s) of interest will continue to be expressed by the transformed mycobacteria following administration of the mycobacteria to a host. Such mycobacteria, therefore, are effective vehicles for expressing antigens which stimulate an immune response, or for the expression of therapeutic(s) agents such as anti-tumor agents and/or anti-cancer agents.

The invention will now be described with respect to the following examples; however, the scope of the present invention is not to be limited thereby.

Example 1

- A. Identification of the DNA sequences of the attachment sites, attB, attL, and attR, of M. smegmatis.

Using standard technologies, a lambda EMBL3 library was prepared from BamHI digested mc²61 chromosomal DNA (mc²61 is a

strain of M. smegmatis which includes an M. smegmatis chromosome into which has been integrated the genome of mycobacterial phage L5) and digested with Bam HI. Phage L5 contains DNA having restriction sites identical to those of phage L1 (Snapper, et al. 1988), except that L5 is able to replicate at 42°C and phage L1 is incapable of such growth. This library was then probed with a 6.7 kb DNA fragment isolated from the L5 genome that had been previously identified as carrying the attP sequence (Snapper, et al 1988). One of the positive clones was plaque purified, DNA prepared, and a 1.1 kb Sal I fragment (containing the AttL sequence) sub-cloned into sequencing vector pUC119. The DNA sequence of this fragment was determined using a shotgun approach coupled with Sanger sequencing. By isolating and sequencing the attL junction site and comparing this to the DNA sequence of L5 that was available, a region was determined where the two sequences aligned but with a specific discontinuity present. The discontinuity represents one side of a core sequence, which is identical in attP, attB, and attL. The region containing the recombinational crossover point is shown in Figure 1.

The attL DNA (1.1 kb Sal I fragment) was used as a probe to hybridize to a Southern blot of Bam HI digested mc²⁶ DNA, which is a strain of M. smegmatis which includes an M. smegmatis chromosome without any phage integration (Jacobs, et al, 1987, hereinabove cited.). A single band of approximately 6.4 kb was detected corresponding to the attB sequence of M. smegmatis. This same attL probe was used to screen a cosmid library of mc²⁶ (provided by Dr. Bill Jacobs of the Albert Einstein College of Medicine of Yeshiva University), and a number of positive cosmid clones were identified. DNA was prepared from these clones, and a 1.9 kb Sal I fragment (containing the attB site) that hybridizes to the attL probe was subcloned into pUC119 for sequencing and further analysis. The DNA sequence containing the core sequence was determined and is shown in Figure 1. The core

sequence, which is identical in attP, attB and attL, has a length of 43bp.

The mc²61 lambda EMBL3 library was then probed with the 1.9kb SalI fragment containing the attB site. Positive plaques were identified, DNA was prepared, and analyzed by restriction analysis and Southern blots. Lambda clones were identified that contained a 3.2kb Bam HI fragment containing the putative attR site. The 3.2kb Bam HI fragment was purified and cloned into pUC119 for sequencing and further analysis.

B. Determination of attP-integrase region of L5 genome.

Concurrent with the above procedures, a significant portion of the DNA sequence of L5 had been determined and represented in several "contigs" or islands of DNA sequence. Sequences of the 6.7kb Bam HI fragment hereinabove described were determined by (a) analysis of the location of Bam HI sites in the contigs of the DNA of L5, and (b) by determining a short stretch of DNA sequence from around the Bam HI sites of plasmid pJR-1 (Figure 6), which carries the 6.7kb Bam HI fragment of L5.

A segment of DNA sequence was located that represented the 6.7kb Bam HI fragment of phage L5. Studies of other phages have shown that the integrase genes are often located close to the attP site. It was thus determined that the L5 integrase (int) gene should lie either within the 6.7kb Bam HI fragment or in a DNA sequence on either side of it. The DNA sequence in the regions was then analyzed by translating it into all six possible reading frames and searching these amino acid sequences for similarity to the family of integrase related proteins, and through computer-assisted analysis of the DNA sequence. As shown in Figure 2, there are shown two domains of reasonably good conservation among L5 integrase and other integrases, and three amino acid residues that are absolutely conserved in domain 2. (See Yagil, et al., J. Mol. Biol., Vol. 207, pgs. 695-717 (1989), and Poyart-Salmeron, et al., J. EMBO, Vol. 8, pgs. 2425-2433 (1989)). A region was identified, and analysis of the

corresponding DNA sequence showed a reading frame that could encode for a protein of approximately 333 amino acids. These observations identified the putative int gene.

The location of the int gene was not within the 6.7kb Bam HI fragment; however, it was very close to it with one of the Bam HI sites (that defines the 6.7kb Bam HI fragment) less than 100 bp upstream of the start of the gene. Analysis of the Bam HI sites showed that the int gene lay within a 1.9kb Bam HI fragment located adjacent to the 6.7kb Bam HI fragment. This 1.9kb Bam HI fragment was cloned by purification of the fragment from a Bam HI digest of L5 DNA and cloning into pUC 119, to generate pMH1 (Figure 7).

From a combination of the above approaches, a schematic of the organization of the attP-int region of L5 was constructed (Figure 3), and the gene sequence of the attP-int region is given in Figure 4.

C. Construction of pMH5.

The 6.7kb Bam HI fragment of mycobacteriophage L5, which contains the attP site, as hereinabove described, was cloned into the Bam HI site of pUC 119 (Figure 5). This was achieved by purifying the 6.7kb Bam HI fragment from a Bam HI digest of L5 DNA separated by agarose gel electrophoresis and ligating with Bam HI cut pUC 119. DNA was prepared from candidate recombinants and characterized by restriction enzyme analysis and gel electrophoresis. A recombinant was identified that contained the 6.7kb Bam HI fragment of L5 cloned into pUC 119. This plasmid was named pJR-1, as shown in Figure 6.

Analysis of DNA sequence data from a project to sequence L5 showed that a 1.9kb Bam HI fragment adjacent to the 6.7kb Bam HI fragment hereinabove described contained the integrase gene.

A plasmid containing a 1.9kb Bam HI fragment containing the DNA encoding for the integrase cloned into the Bam HI site of pUC 119 was constructed. The 1.9kb fragment was purified from a Bam

HI digest of L5 DNA and cloned into the Bam HI site of pUC 119. Construction of the recombinant was determined by restriction analysis and gel electrophoresis. This plasmid was called pMH1, the construction of which is shown schematically in Figure 7.

pJR-1 was then modified by digestion with EcoRI and SnaBI (both are unique cloning sites), between which is a Bam HI site. The Eco RI-Sna BI fragment, including the Bam HI site was excised, and the plasmid was religated to form plasmid of pMH2, which contains one Bam HI site compared to two Bam HI sites contained in pJR-1. A schematic of the construction of pMH2 is shown in Figure 8.

The 1.9kb Bam HI fragment, which includes the integrase gene, was purified from a Bam HI digest of pMH1 and ligated to Bam HI digested pMH2. Recombinants were identified as above and the orientation of the 1.9kb fragment determined. A plasmid called pMH4 was thus constructed (Figure 9) in which the region from the Sna BI site (upstream of attP) through to the Bam HI site (downstream of the integrase gene) was identical to that in L5.

pMH4 was digested with HindIII (unique site) and was ligated to a 1kb HindIII fragment purified from pKD43 (supplied by Keith Darbyshire of the Nigel Gindley Laboratory) that contains the gene determining resistance to kanamycin. Recombinants were identified and characterized as above. This plasmid is called pMH5. A schematic of the construction of pMH5 is shown in Figure 10.

D. Integration of pMH5 into attB of *M. smegmatis*.

Plasmids pYUB12 (a gift from Dr. Bill Jacobs a schematic of the formation of which is shown in Figure 20), pMD01 (Figure 11), and pMH5 were electroporated, with four different concentrations of plasmid DNA over a 1,000-fold range, into *M. smegmatis* strain mc²155, a strain which is able to support plasmid replication. In Examples 1 and 2, all electroporation procedures of *M. smegmatis*, or of BCG, were carried out as follows:

Cultures of organism were grown in Middlebrook 7H9 media, as described by Snapper, et al. (1988), harvested by centrifugation, washed three times with cold 10% glycerol, and resuspended at approximately a 100 x concentration of cells.

1 μ l of DNA was added to 100 μ l of cells in an ice-cold cuvette and pulsed in a Bio-Rad Gene Pulser, and given a single pulse at 1.25 kv at 25 μ F. 1 ml of broth was added the cells incubated for 1 hr. at 37°C for expression of the antibiotic-resistant marker. Cells were then concentrated and plated out on Middlebrook or tryptic soy media containing 15 μ g/ml kanamycin. Colonies were observed after 3 to 5 days incubation at 37°C.

Each of pYUB12, pMD01, and pMH5 carries kanamycin resistance. Plasmid pYUB12 carries an origin of DNA replication, while pMD01 lacks a mycobacterial origin of replication. Plasmid pMH5 does not carry a mycobacterial origin of replication, but carries a 2kb region of phage L5 which contains the attP site and the integrase gene (Figure 4). The number of transformants were linear with DNA concentration. Plasmid pYUB12 gives a large number of transformants (2×10^5 per μ g DNA) in mc²155, while pMH5 gives 6×10^4 transformants per μ g DNA, and pMD01 gives no transformants.

The above experiment was then repeated by electroporating the plasmids pYUB12, pMD01, and pMH5 into M. smegmatis strain mc²6, which does not support plasmid replication. No transformants in mc²6 were obtained from pYUB12 or pMD01, while pMH5 gave approximately 10^4 kanamycin resistant transformants in mc²6 per μ g of DNA, thus indicating integration of pMH5 into the mc²6 chromosome.

DNA from six independent pMH5 transformants (four in mc²155 and two in mc²6) was prepared. These DNA's (along with DNA from both mc²155 itself, and mc²155 carrying the plasmid pYUB12) were digested with a restriction enzyme, and analyzed by Southern blot and hybridization with the M. smegmatis 1.9kb attB probe

hereinabove described. As shown in Figure 12, all six transformants have integrated into the attB site, resulting in the production of two new DNA fragments with different mobilities. If pMH5 did not integrate into the attB site, it would be expected that a single band, corresponding to the attB site in the mc²155 control, would be obtained.

E. Construction of pMH9.2 and pMH9.4

pUC119 was digested with HindIII, and a 1kb HindIII fragment, containing a kanamycin resistance gene, purified from pKD43, was ligated to the HindIII digested pUC119 to form pMH8 (Figure 13.). A 2kb SalI fragment of pMH5 (bp 3226-5310), which carries the attP and integrase gene from SalI digested pMH5, was purified and inserted in both orientations relative to the vector backbone of SalI digested pMH8 to form plasmids pMH9.2 and pMH9.4 (Figures 14 and 15).

F. Stability of plasmids in M. smegmatis

M. smegmatis strain mc²155 cells carrying, as a result of electroporation, plasmid pYUB12, pMH9.2 or pMH9.4, or strain mc²6 cells carrying plasmid pMH5, as a result of electroporation as hereinabove described, were grown to saturation in broth with kanamycin. Cultures were then diluted 1:100 into broth without kanamycin and grown to saturation. Two further cycles of dilution and growth were done, corresponding to about 20 generations of bacterial growth. Cultures were plated out to single colonies on non-selective plates, and approximately 100 of these colonies were patch plated onto both non-selective and selective plates. The % of colonies that were sensitive to kanamycin, thus corresponding to the percentage of cells which lost the plasmid, is given below in Table I.

Table I

	<u>% loss</u>
pYUB12 (mc ² 155)	35
pMH5 (mc ² 6)	17
pMH9.2 (mc ² 155)	3

pMH9.4 (mc²155)

O

Example 2

The 1.9 kb Sal I fragment, which includes the M. smegmatis attB site as hereinabove described was cloned into pUC119, and the plasmid generated was named pMH-12. (Figure 16).

Gel purified Sal I 1.9kb M. smegmatis fragment containing attB (isolated from pMH-12) was used to probe a Southern transfer of Bam HI digested mycobacterial DNA's, including BCG substrain Pasteur, shown in Figure 17. This demonstrated that there is one Bam HI fragment of BCG that strongly hybridizes to the M. smegmatis attB probe and three hybridize weakly. The strongest hybridizing band is the fastest moving band (approximately 1.9 kb).

The same probe as above was used to probe a BCG cosmid library (provided by Dr. Bill Jacobs) and positive clones were identified. DNA was prepared from several positive clones and analyzed by restriction analysis and Southern blotting. The 1.9 kb Bam HI fragment (corresponding to the strongly hybridizing band in the Southern blot) was identified, gel purified from the cosmid DNA and cloned into pUC119. The resulting plasmid was named pMH-15. (Figure 18).

Plasmids pMH-5 and pMH9.4 were electroporated into BCG Pasteur. It was observed that pMH9.4 transforms BCG with high efficiency (approximately 10^4 transformants/ μ g DNA), while pMH-5 transforms BCG at low efficiency (1-10 transformants/ μ g DNA). DNA was prepared from BCG transformants and analyzed by Bam HI restriction and Southern blot analysis, probing with gel purified 1.9kb Bam HI BCG attB fragment from pMH-15. These data are shown in Figure 19 and show that integration of both pMH 5 and pMH 9.4 is specific to the BCG attB site (ie. the strongly cross-hybridizing fragment in BCG). This is illustrated by the loss of the 1.9kb Bam HI fragment from the transformants and the appearance of two new bands representing attL and attR junction fragments. Figure 19 shows just one of the pMH 5/BCG

transformants, although all of the four that were analyzed show that one of the bands (the largest) is smaller than expected (and different in each of the transformants), indicating that the transformation efficiency of pMH 5 is low in BCG. In contrast, the four pMH 9.4 transformants are identical to each other (Figure 19) and give attR and attL junction fragments of the predicted sizes.

Example 3

A. Construction of plasmids including mycobacterial promoter expression cassette.

1. Construction of pYUB125

Plasmid pAL5000, a plasmid which contains an origin of replication of M. fortuitum, and described in Labidi, et al., FEMS Microbiol. Lett., Vol. 30, pgs. 221-225 (1985) and in Gene, Vol. 71, pgs. 315-321 (1988), is subjected to a partial Sau 3A digest, and 5kb fragments are gel purified. A 5kb fragment is then ligated to Bam HI digested pIJ666 (an. E. coli vector containing an E. coli origin of replication and also carries neomycin-kanamycin resistance, as described in Kieser, et al., Gene, Vol. 65, pgs. 83-91 (1988) to form plasmid pYUB12. A schematic of the formation of plasmid pYUB12 is shown in Figure 20. pYUB12 and pIJ666 were then transformed into M. smegmatis and BCG. Neomycin-resistant transformants that were only obtained by pYUB12 transformation confirmed that pAL5000 conferred autonomous replication to pIJ666 in M. smegmatis and BCG.

Shotgun mutagenesis by Snapper, et al (1988, hereinabove cited) indicated that no more than half of the pAL5000 plasmid was necessary to support plasmid replication in BCG. This segment presumably carried open reading frames ORF1 and ORF2, identified by Rauzier, et al., Gene, Vol. 71, pgs. 315-321 (1988), and also presumably carried a mycobacterial origin of replication. pYUB12 is then digested with HpaI and EcoRV, a 2586 bp carrying this region or segment of pAL5000 is removed and

ligated to PvuII digested pYUB8. Plasmid pYUB8 (a pBR322 derivative) includes an E. coli replicon and a kan^R (aph) gene. Ligation of the 2586 bp pYUB12 fragment to PvuII digested pYUB8 results in the formation of pYUB53, as depicted in Figure 21. Transformation of pYUB53 confirmed that the EcoRV-HpaI fragment, designated M.rep, was capable of supporting autonomous replication in BCG.

Plasmid pYUB53 was then digested with AatI, EcoRV, and PstI in order to remove the following restriction sites:

AatI 5707
EcoRI 5783
BamHI 5791
Sali 5797
PstI 5803
PstI 7252
Sali 7258
BamHI 7264
EcoRI 7273
ClaI 7298
HindIII 7304; and
EcoRV 7460

Fragment ends are then flushed with T4 DNA polymerase and religated to form plasmid pYUB125, construction of which is shown in Figure 22.

2. Elimination of superfluous vector DNA from pYUB125

792 bases of the tet gene, which had been inactivated by prior manipulations, was eliminated by a complete NarI digest, gel purification of the 6407 bp fragment, and ligation/recirculation, transformation of E. coli strain HB101, and selection of Kan^R transformants. The construction of the resulting plasmid, pMV101, is schematically indicated in Figure 23, and the DNA sequence of pMV101, which includes markings of regions which will be deleted and of mutations, as hereinafter described, is shown in Figure 24.

3. Elimination of undesirable restriction sites in aph (kan^R) gene.

To facilitate future manipulations, the HindIII and ClaI restriction sites in the aph gene were mutagenized simultaneously by polymerase chain reaction PCR mutagenesis according to the procedure described in Gene, Vol. 77 pgs. 57-59 (1989). The bases changed in the aph gene were at the third position of codons (wobble bases) within each restriction site and the base substitutions made were designed not to change the amino acid sequence of the encoded protein.

Separate PCR reactions of plasmid pMV101 with primers ClaMut-Kan + HindRMut-Kan and HindFMat-Kan + Bam-Kan were performed at 94°C (1 min.), 50°C (1 min.), and 72°C (1 min.) for 25 cycles. The PCR primers had the following base sequences:

ClaMut-Kan

CTT GTA TGG GAA GCC CC

HindRMut-Kan

GTG AGA ATG GCA AAA GAT TAT GCA TTT CTT TCC AG

HindFMat-Kan

GTC TGG AAA GAA ATG CAT AAT CTT TTG CCA TTC TCA CCG G

Bam-Kan

CGT AGA GGA TCC ACA GGA CG

The resulting PCR products were gel purified and mixed and a single PCR reaction without primers was performed at 94°C (1 min.), 72°C (1 min.) for 10 cycles. Primers ClaMut-Kan and Bam-Kan were added and PCR was resumed at 94°C (1 min.), 50°C (1 min.), and 72°C (2 min.) for 20 cycles. The resulting PCR product (Kan. mut) was digested with BamHI and gel purified. Plasmid pMV101 was digested with ClaI and cohesive ends were filled in by Klenow + dCTP + dGTP. Klenow was heat inactivated and the digest was further digested with BamHI. The 5232 base pair fragment was gel purified and mixed with fragment Kan.mut and ligated. The ligation was transformed into E.coli strain HB101 and Kan^R colonies were screened for plasmids resistant to

ClaI and HindIII digestion. Such plasmids were designated as pMV110, which is depicted in Figure 23.

4. Elimination of sequences not necessary for plasmid replication in mycobacteria.

Plasmid pMV110 was resected in separate constructions to yield plasmids pMV111 and pMV112. In one construction, pMV110 was digested with NarI and Ball, the ends were filled in, and a 5296 base pair fragment was ligated and recircularized to form pMV111. In another construct, pMV110 was digested with NdeI and SphI, the ends were filled in, and a 5763 base pair fragment was ligated and recircularized to form pMV112. Schematics of the constructions of pMV111 and pMV112 are shown in Figure 25. These constructions further eliminated superfluous E. coli vector sequences derived from pAL5000 not necessary for mycobacterial replication. Cloning was performed in E. coli. Plasmids pMV111 and pMV112 were tested for the ability to replicate in M. smegmatis. Because both plasmids replicated in M. smegmatis the deletions of each plasmid were combined to construct pMV113.

To construct pMV113 (Figure 25), pMV111 was digested with BamHI and EcoRI, and a 1071 bp fragment was isolated. pMV112 was digested with BamHI and EcoRI, and a 3570 bp fragment was isolated, and then ligated to the 1071 bp fragment obtained from pMV111 to form pMV113. These constructions thus defined the region of pAL5000 necessary for autonomous replication in mycobacteria as no larger than 1910 base pairs.

5. Mutagenesis of restriction sites in mycobacterial replicon.

To facilitate further manipulations of the mycobacterial replicon, PCR mutagenesis was performed as above to eliminate the Sal I, EcoRI, and BglII sites located in the open reading frame known as ORF1 of pAL5000. PCR mutagenesis was performed at wobble bases within each restriction site and the base substitutions were designed not to change the amino acid sequence of the putative encoded ORF1 protein. The restriction sites were eliminated one at a time for testing in mycobacteria. It was

possible to eliminate the SalI and EcoRI without altering replication in M. smegmatis. In one construction, PCR mutagenesis was performed at EcoRI1071 of pMV113 with primers Eco Mut - M.rep and Bam-M.rep. to form pMV117, which lacks the EcoRI1071 site. Primer Eco Mut - M.rep has the following sequence:

TCC GTG CAA CGA CGT GTG TCC CGG A;

and Bam-M.rep has the following sequence:

CAC CCG TCC TGT GGA TCC TCT AC.

In another construction, PCR mutagenesis was performed at the SalI 1389 site with primer Sal Mut - M.rep and Bam-M.rep to form pMV119, which lacks the SalI 1389 site. Primer Sal Mut - M.rep has the following sequence:

TGG CGA CCG CAG TTA CTC AGG CCT.

pMV117 was then digested with ApaLI and BglII, and a 3360 bp fragment was isolated. pMV119 was digested with ApaLI and BglII, and a 1281 bp fragment was isolated and ligated to the 3360 bp fragment isolated from pMV117 to form pMV123. A schematic of the constructions of plasmids pMV117, pMV119, and pMV123 is shown in Figure 26. Elimination of the BglII site, however, either by PCR mutagenesis or Klenow fill in, eliminated plasmid replication in mycobacteria, thus suggesting that the BglII site is in proximity to, or within a sequence necessary for mycobacteria plasmid replication.

6. Construction of pMV200 series vectors.

To facilitate manipulations of all the components necessary for plasmid replication in E. coli and mycobacteria, (E. rep. and M. rep.) and selection of recombinants (Kan^R), cassettes of each component were constructed for simplified assembly in future vectors and to include a multiple cloning site (MCS) containing unique restriction sites and transcription and translation terminators. The cassettes were constructed to allow directional cloning and assembly into a plasmid where all transcription is unidirectional.

Kan^R Cassette

A DNA cassette containing the aph (Kan^R) gene was constructed by PCR with primers Kan5' and Kan3'. An SpeI site was added to the 5' end of PCR primer Kan3', resulting in the formation of a PCR primer having the following sequence:

CTC GAC TAG TGA GGT CTG CCT CGT GAA G.

Bam HI + NheI sites were added to the 5' end of PCR primer Kan5', resulting in the formation of a PCR primer having the following sequence:

CAG AGG ATC CTT AGC TAG CCA CCT GAC GTC GGG G.

PCR was performed at bases 3375 and 4585 of pMV123, and BamHI and NheI sites were added at base 3159, and an SpeI site was added at base 4585. Digestion with BamHI and SpeI, followed by purification resulted in a 1228/2443 Kan^R cassette bounded by BamHI and SpeI cohesive ends with the direction of transcription for the aph gene proceeding from BamHI to Spe I.

E. rep. cassette

A DNA cassette containing the ColEI replicon of pUC19 was constructed by PCR with primers E.rep/Spe and E.rep/Mlu. An SpeI site was added to the 5' end of PCR primer E.rep/Spe, and an MluI site was added to the 5' end of PCR primer E.rep./Mlu. The resulting primers had the following base sequences:

E.rep./Spe

CCA CTA GTT CCA CTG AGC GTC AGA CCC

E.rep./Mlu

GAC AAC GCG TTG CGC TCG GTC GTT CGG CTG.

PCR was performed at bases 713 and 1500 of pUC19, and an MluI site was added to base 713, and a SpeI site was added to base 1500. Digestion with MluI and SpeI, followed by purification resulted in an E.rep. cassette bounded by SpeI and MluI cohesive ends with the direction of transcription for RNA I and RNA II replication primers proceeding from SpeI to MluI.

M.rep. cassette

A DNA cassette containing sequences necessary for plasmid replication in mycobacteria was constructed by PCR of pMV123 with primers M.rep/Mlu and M.rep/Bam. An MluI site was added to the 5' end of PCR primer M.rep/Mlu. A BamHI site was added to the 5' end of PCR primer M.rep/Bam. The resulting PCR primers had the following base sequences:

M.rep./Mlu

CCA TAC GCG TGA GCC CAC CAG CTC CG

M.rep./Bam

CAC CCG TCC TGT GGA TCC TCT AC

PCR was performed at bases 134 and 2082 of pMV123. An MluI site was added to base 2082. Digestion with BamHI and MluI, followed by gel purification resulted in a 1935 base pair DNA cassette bounded by MluI and BamHI cohesive ends with the direction of transcription for the pAL5000 ORF1 and ORF2 genes proceeding from MluI to Bam HI.

The Kan^R, E.rep, and M.rep PCR cassettes were then mixed in equimolar concentrations and ligated, and then transformed in E.coli strain HB101 for selection of Kan^R transformants. Colonies were screened for the presence of plasmids carrying all three cassettes after digestion with BamHI + MluI + SpeI and designated pMV200. An additional restriction site, NcoI, was eliminated from the M.rep cassette by digestion of pMV200 with NcoI, fill in with Klenow, and ligation and recircularization, resulting in the formation of pMV201. A schematic of the formation of pMV200 from pMV123 and pUC19, and of pMV201 from pMV200, is shown in Figure 27. Plasmids pMV200 and pMV201 were transformed into M. smegmatis and BCG. Both plasmids yielded Kan^R transformations, thus indicating their ability to replicate in mycobacteria.

A synthetic multiple cloning sequence (MCS) (Figure 28) was then designed and synthesized to facilitate versatile molecular cloning and manipulations for foreign gene expressions in mycobacteria, and for integration into the mycobacterial

chromosome. The synthetic MCS, shown in Figure 28, contains 16 restriction sites unique to pMV201 and includes a region carrying translation stop codons in each of three reading frames, and a transcription terminator derived from *E. coli* 5S ribosomal RNA (T1).

To insert the MCS cassette, pMV201 was digested with *NarI* and *NheI*, and the resulting fragment was gel purified. The MCS was digested with *HinPI* and *NheI* and, the resulting fragment was gel purified. The two fragments were then ligated to yield pMV204. A schematic of the construction of pMV204 is shown in Figure 29.

Plasmid pMV204 was then further manipulated to facilitate removal of the M.rep cassette in further constructions. pMV204 was digested with *MluI*, and an *MluI* - *Not I* linker was inserted into the *MluI* site between the M.rep and the E.rep to generate pMV206. A schematic of the construction of pMV206 from pMV204 is shown in Figure 30, and the DNA sequence of pMV206 is given in Figure 31.

7. Construction of expression cassette based on BCG HSP60.

Among the most abundant proteins in mycobacteria is the HSP60 heat shock protein (also known as the 65 kda antigen). Because abundance of the HSP60 protein in mycobacteria indicates strong HSP60 gene expression, the sequence controlling HSP60 expression was chosen to control expression of heterologous genes encoding antigens or other proteins in BCG.

The published sequence of the BCG HSP60 gene (Thole, et al, Infect. and Immun., Vol. 55, pgs. 1466-1475 (June 1987)), and surrounding sequence permitted the construction of a cassette carrying expression control sequences (i.e., promoter and translation initiation sequences) by PCR. The BCG HSP61 cassette (Figure 32) contains 375 bases 5' to the BCG HSP60 start codon, and 15 bases (5 codons) 3' to the start codon. PCR oligonucleotide primers were then synthesized. Primer Xba-HSP60, of the following sequence:

CAG ATC TAG ACG GTG ACC ACA ACG CGC C
was synthesized for the 5' end of the cassette, and primer Bam-HSP61, of the following sequence:

CTA GGG ATC CGC AAT TGT CTT GGC CAT TG
was synthesized for the 3' end of the cassette. The primers were used to amplify the cassette by PCR from BCG substrain Pasteur chromosomal DNA. The addition of the Bam HI site at the 3' end of the cassette adds one codon (Asp) to the first six codons of the HSP60 gene.

Each of pMV206 and the PCR cassette HSP61 was digested with Xba I and Bam HI. The PCR cassette was then inserted between the Xba I and Bam HI sites of pMV206, and then ligated to form pMV261, which is shown in Figure 33.

The E. coli lac Z gene was used as a reporter, or marker gene to assay the ability of the HSP61 cassette to express heterologous genes in BCG. A BamHI restriction fragment carrying the lac Z gene was cloned into the Bam HI site of Bam HI digested pMV261, resulting in the formation of pMV261/LZ. A schematic of the construction of pMV261/LZ is shown in Figure 34. The formation of pMV261/LZ results in a fusion between the HSP60 and lac Z genes at the sixth codon of the HSP60 gene and the sixth codon of the lac Z gene. pMV261/LZ was then transformed into E. coli. Blue E. coli colonies were selected on x-gal plates for the presence of pMV261/LZ, thus indicating that the HSP60 promoter and translation initiation sequences were also active in E. coli.

pMV261/LZ was then transformed into BCG and plated on Dubos Oleic Agar plates containing x-gal. All BCG colonies resulting from this transformation exhibited blue color, thus indicating that the lac Z gene product (B-galactosidase) was expressed in BCG. SDS polyacrylamide gel electrophoresis was performed on lysates of the pMV261/LZ BCG recombinant, revealing that B-galactosidase protein was expressed to levels in excess of 10% of total BCG protein (as determined by staining with Coomassie

brilliant blue). These data indicated that BCG HSP61 expression cassette was functional in expression vector pMV261, and that substantial expression of a heterologous gene could be achieved using HSP60 - controlled expression in BCG.

Plasmid pMV261/LZ was then shown to replicate autonomously, and express the E. coli B-galactosidase, or lacZ gene, driven by the BCG promoter HSP60, in M. smegmatis and BCG.

B. Transfer of mycobacterial phage L5 integration sequences to BCG expression vector.

Plasmid pMH9.4, which includes the mycobacterial phage L5 attP site and the L5 integrase gene, was digested to completion with either KpnI + PvuII or XbaI + PvuII, and a restriction fragment of 1862 or 1847 base pairs, respectively, each of which contain the attP site and the integrase gene, were purified by agarose gel electrophoresis. Plasmid pMV261/LZ was digested with XbaI + DraI to generate either a 7569 bp or 7574 bp vector fragment. The 7569 bp fragment was ligated to the 1862 bp fragment derived from pMH9.4 to form pMV460F/LZ. The 7574 bp fragment was ligated to the 1847 bp fragment derived from pMH9.4 to form pMV460 R/LZ. Plasmids pMV460 F/LZ and pMV460R/LZ each include a mycobacterial replicon, the L5 attP site, and the L5 integrase gene. A schematic of the formation of plasmids pMV460 F/LZ and pMV460R/LZ is shown in Figure 35. To generate derivatives without the mycobacterial plasmid replicon, plasmids pMV460F/LZ and pMV460R/LZ were digested with NotI and recircularized by ligation to generate pMV360F/LZ and pMV360R/LZ. A schematic of the construction of pMV360F/LZ and pMV360R/LZ is shown in Figure 36.

Plasmids pMH9.4, pMV261/LZ, pMV460F/LZ, pMV460R/LZ, pMV360F/LZ, and pMV360R/LZ were then transformed into M. smegmatis and BCG to test their ability to replicate autonomously or integrate into the M. smegmatis or BCG chromosome. Transformation with pMH9.4, pMV261/LZ, pMV360F/LZ, and pMV360R/LZ yielded kanamycin resistant transformants of M. smegmatis and

BCG. Transformants of pMV261LZ, pMV360F/LZ, and pMV360R/LZ were shown to express E. coli B-galactosidase by SDS-polyacrylamide gel electrophoresis and X-gal assay as hereinabove described for pMV261/LZ. Plasmids pMV461F/LZ and pMV461R/LZ failed to yield kanamycin resistant transformants, thus indicating that chromosomal integration of a plasmid carrying sequences mediating autonomous replication is lethal to mycobacteria.

Example 4

A. Construction of expression cassettes based on BCG HSP70.

A partial sequence of the 5' region of the BCG HSP70 gene (which encodes for the BCG HSP70 heat shock protein, also known as the 70 kda antigen) obtained by Dr. Rick Young (MIT) permitted the construction of cassettes carrying expression control sequences (i.e., promoters and translation initiation sequences) by PCR, according to the procedures hereinabove cited. The BCG-HSP71 cassette (Figure 32) contains 150 bases 5' to the BCG-HSP70 start codon and 15 bases (5 codons) 3' to the start codon. Primer Xba-HSP70 was synthesized for the 5' end of the cassette, and primer Bam-HSP71 was synthesized for the 3' end of the cassette. The primers had the following base sequences:

Xba-HSP70

GGC CTC TAG ACC CGC ACG ACC AGC GTT AGC

Bam-HSP71

GCT AGG ATC CCC GAC CGC ACG AGC CAT GGT

The primers were used to amplify the cassette from BCG substrain Pasteur chromosomal DNA. The addition of the Bam HI site at the 3' end of the cassette adds 1 codon (Asp) to the 3' end of the HSP71 expression cassette.

Each of pMV206 and the PCR cassette HSP71 was digested with XbaI and BamHI. The PCR cassette was then inserted between the XbaI and BamHI sites of PMV 206, and then ligated to form pMV271, which is shown in Figure 33.

B. Cloning of HIV-1gag.

A BamHI-ClaI PCR cassette of HIV-1 gag was cloned between the Bam HI and Cla I sites of pMV261 and pMV271 to obtain pMV261/gag and pMV271/gag. Expression of the gag antigens in BCG was verified by the appearance of immunoreactive protein bands in Western blot analysis of BCG pMV271/gag recombinant lysates. BCG transformants, however, were never obtained with pMV261/gag, thus indicating that gag as expressed from pMV26/gag was lethal.

C. Integration of HSP60-gag expression cassette into BCG.

In order to test whether integration of an HSP-60-gag expression cassette into BCG would result in non-lethal expression of gag in BCG, it was decided that the HSP60-gag expression cassette be cloned into a plasmid (pMV307) which includes the mycobacteriophage L5 attP and integrase sequences, the construction of which is explained hereinbelow.

1. Construction of pMV307.

Plasmid pMV206 was digested with NotI to remove the mycobacterial replicon. The resulting 2209 bp fragment, which includes the aph (Kan^R) gene, the E.coli replicon and the multiple cloning site, was ligated and recircularized to form pMV205, the construction of which is schematically depicted in Figure 30.

PCR with primers XbaI-Att/Int and NheI-Att/Int was then performed on a Sal I fragment from pMH9.4, which contains the attP site and the L5 integrase gene. The resulting cassette was then digested with XbaI and NheI, and a 1789 bp fragment was gel purified. pMV205 was then digested with NheI, and the resulting fragment was ligated to the 1989 bp fragment obtained from pMH9.4 to form pMV307. A schematic of the construction of pMV307 is shown in Figure 37.

2. Construction and transformation of pMV361/gag.

The XbaI-ClaI HSP-antigen cassette, which includes the HSP60 promoter and HIV-1gag sequences, was cloned between the NotI and ClaI sites of pMV307 to form plasmid pMV361/gag. pMV361/gag was

then transformed into BCG and shown to express HIV-1 gag protein by Western blot analysis with HIV-1 infected human sera.

Example 5

A series of antigen gene fragments, or cassettes, were constructed by PCR, with the exception of the gene fragment containing the gene for human tumor antigen p97, as indicated in Table 2, and cloned into various restriction sites of pMV261 and pMV271 to form new constructs of the pMV261 and pMV271 series. The antigen genes, antigen gene fragments, cloning sites used in pMV261 and pMV271 and the names of the resulting constructs, are given below in Table 2.

The following constructs:

pMV261/P97
pMV271/P97
pMV261/CS
pMV271/CS
pMV261/SM97
pMV271/SM97
pMV271/HIV-1-gag
pMV261/HIV-1gp120
pMV271/HIV-1gp120
pMV261/HIV-1-gp41
pMV271/HIV-1-gp41
pMV261/HIV2gag
pMV271/HIV2gag
pMV261/HIV2-gp120
pMV271/HIV2-gp120
pMV261/HIV2-gp41; and
pMV271/HIV2-gp41.

were transformed into BCG, and the presence of the corresponding antigens in BCG was verified by the appearance of immunoreactive protein bands in Western blot analysis of BCG recombinant lysates.

Example 6

Antigen gene expression cassettes, which include a promoter sequence and a heterologous gene sequence, were excised from the pMV261 and pMV271 derivatives with NotI and a second restriction enzyme site (Pvu II, Eco RI, Sal I, Cla I or Hind III) and cloned into the integrating plasmid pMV307 between the NotI site and a second enzyme site (Pvu II, Eco RI, Sal I, Cla I or Hind III) to form the pMV 361/XXX and pMV371/XXX series of plasmids (e.g., pMV361/HIV-1gp120). The backbones of these series of plasmids (pMV361 and pMV371) are shown in Figure 38.

The following plasmids:

pMV361/P97

pMV371/P97
pMV361/CS
pMV371/CS
pMV361/SM97
pMV371/SM97
pMV361/HIV-1gag
pMV371/HIV-1gag
pMV361/HIV-1gp120
pMV371/HIV-1gp120
pMV361/HIV-1gp41
pMV371/HIV-1gp41
pMV361/HIV2gag
pMV371/HIV2gag
pMV361/HIV2-gp120; and
pMV361/HIV2-gp41.

were transformed into BCG and shown to express the corresponding antigens by Western blot analysis with the appropriate antigen-specific human sera.

It is to be understood, however, that the scope of the present invention is not to be limited to the specific embodiments described above. The invention may be practiced other than as particularly described and still be within the scope of the accompanying claims.

WHAT IS CLAIMED IS:

1. A DNA for integrating DNA into a mycobacterium chromosome, comprising:
a first DNA sequence which is a phage DNA portion encoding bacteriophage integration into a mycobacterium chromosome; and
a second DNA sequence encoding a protein or polypeptide which is heterologous to the mycobacterium in which the DNA is to be integrated.
2. The DNA of Claim 1 wherein said phage DNA portion encodes mycobacteriophage integration into a mycobacterium chromosome.
3. The DNA of Claim 1 wherein said first DNA sequence includes DNA encoding integrase.
4. The DNA of Claim 3 wherein said first DNA sequence includes DNA encoding an AttP site.
5. The DNA of Claim 4 wherein said first DNA sequence encodes integration into BCG.
6. The DNA of Claim 5 wherein the DNA is a plasmid including an origin of replication for E. coli.
7. The DNA of Claim 1 wherein the first DNA sequence is selected from the group consisting of Figure 4 and analogues and derivatives thereof which encode for integration into mycobacteria.
8. Mycobacteria transformed with the DNA of Claim 1.
9. BCG having foreign DNA integrated into the chromosome.
10. A vaccine comprising: the transformed mycobacteria of Claim 8; and an acceptable pharmaceutical carrier.
11. A process for providing a protein, comprising:
expressing the protein in the mycobacteria of Claim 8.

AAAACACCCCTCTGACCAGCCGAGCGGGCGACGGGAATCGAACCCGCGTAGCTAGTTTGGGAAGATGGGTGTCTGCCGACCACA attP

AAAACACCCCTCTGACCAGCCGAGCGGGCGACGGGAATCGAACCCGCGTAGCTAGTTTGGGAAGACTAGGGCTCTACCATTGAGC attL

CGCACCGTGGCGGTCCCTACCGAGCGGGCGACGGGAATCGAACCCGCGTAGCTAGTTTGGGAAGACTAGGGCTCTACCATTGAGC attB

Figure 1

Domain 1

INT (λ) RLAHELAVVTGQGVGDLCEMKKWSDIVDG
 INT (HK22) RLANDLAVVTGQGVGDLCEMKKWSINDN
 INT (φ 80) VFLVKEIMLTGCRTAEIRLSERSWFRLD
 INT (P2) KKIAILCLSTGARWGEAARLKAENI IHN
 INT (P4) MIAVKLSLLTFVRSSELRFARWDEFDD
 INT (P22) KSVVEFALSTGLRNSNIINLEMQQIDMQ
 INT (186) ETVVRICLATGARWSEAESLAKSQLAKY
 INT (HP1) GLIVRICLATGARWSENETLTQSQVMPY
 INT (L54a) AGAVEQALTCMRIGELLALQVKDVLK
 Cre (P1) TAGVEKALSGLVTKLVERWISVSGVADD
 D Prot. (F) KMLLATLWNTGARINEALALTRGDFSLA
 Flm B YCLTLICFINGFRASEICRLRISDIDLK
 Flm E YCLILLAYRHGMRISELDDLHYQDIDLN
 Tn2603 ORF3 RLFAQLLYGTGMRISEGLQLRVKDIDED
 Tn554 TnpA KLIIMHYEGGLRIGEVLSIRLEDIVTN
 Tn554 TnpB ATHTHIVQECGMRISELCTLRKKGCLIED
 Tn4430 TnpI YAIATLLAYTGVRISEALSIRKNDENLQ
 Rci YVIFILALETAMRQOEILALRWEHIDLR
 Tn1545 ORF2 YDEILILLKTGLRISSEFGGLTLPLDIDFE
 Flp -----

CONSENSUS --lv-l-l-TGmR-SEl--Lr---di---

L5 RIANYILANTSLRFGELIELRRKDIVDD

Domain 2

HELRLSLA-RLYEKQ-ISDKFAQHLLGHIKS-DTHASQYR-
 HELRLSLA-RLYRNQ-IGDKFAQRLGHIKS-DSMAARYRD
 HDHRTIATNLSELG-CPPHVIEKLLGHQM-VGVMAHYN-
 HIALRHSFATHFING-GSIIILQRIILGHTR-IEQTMVYAH
 HGFRTHMARGALGESGLWSODAJERQSLHSENNVRAAYIH
 HDLRITWASHLVQAG-VPISVLQEMGGWES-IEHVRRYAH
 HVLRLITFASHFENG-GNILLVQLQVGLHTD-IRKMTTRYAH
 HVLRLITFASHFENG-GNILLVKEILGHST-IEMTTRYAH
 HTLRHTHISLLAEMN-ISLKAIMKRVGHRDEKTIKVVYTH
 HSARVGAARDMARAG-VSIEIMOAGGWTN-VNIVMNYIR
 HTFRHSYAMHMLYAG-IPLKVLQSLMGHS-ISSTEVTYK
 HMLRHSCGFALANMG-IDTRLIQDYLGHRN-IRHTVRYTA
 HMLRHACGYELAERG-ADTRLIQDYLGHRN-IRHTVRYTA
 HTLRHSFATALLRSG-YDINTVQDLLGHSD-VSTTHIYTH
 HMLRHTHATQLIREG-WDVAFVQKRLGHANVQTTLTIVH
 HAFRUITVGTIRMING-HPQHIQVQKFLGHES-PEHTSRYAH
 HQLRHUFFCTNAIEKG-FSIEHEVANQAGHSN-IHTTLTYT-
 HDLRHEAISRFEELGSLAVVHEIAANISGHS-MNMLKRYTH
 HSLRHTFCTNYANAG-MNPKALQYINGHAN-IAHTLNYIA
 HIGRHLMTSFLSMKGLTELTVVGVNWSDKRASAVATTYTH

* *
 H-LRht-At-l---G---l--lQ-lLgh---l---T---Y-H

HDLRVAGATFAAQAG-ATTKELMARLGHST-PRMANKYQM

Figure 2

L5 Bam HI sites





Figure 3

Figure 4

1301 GACCGTCCGGTCCGAAGGGTCTGTGACGGTTCCGGCTCAGGTCCGGGAGATGATCCGAGCGCACATGAAGGACCGTACGAAGATGAACAAAGGGCCCGAG
Int CTGGCAGGCCAGCTTCGCAGGACACTGCCAAGGGCGAGTCCAGCGCTCTAGTAGGCTCGCGTGTACTTCTTGGCATGCTTCTACTGTTCGGGGCTC 1400
T V R S K R P V T V P P H V A E M I R A H M K D R T K M N K G P E -

1401 GCATTCTGGTGACCACGACGCGAGGGCAACCGGCTGTGGAAGTCCGGCTTCACCAAGTCGCTGAAGCGTGGCTACGGCAAGATCGGTGCGCGGAACTCC
Int CGTAAGGACCACTGGTGTGCGTCCCGTTGGCGGACAGCTTCAGGGCGAAGTGGTTCAGCGACTTCGCACCGATGCGGTTCTAGCCAGCGGGCTTGAGG 1500
A F L V T T T Q G N R L S K S A F T K S L K R G Y A K I G R P E L R -

1501 GCATCCACGACCTCCGCGCTGTGCGGCTACGTTCCGGCTCAGGCAAGGTGCGACGACCAAGGAGGTGATGGCGCGTCTCGGTACACGACTCTTAGGAT
Int CGTAGGTCTGGAGGCGCGACAGCGCGATGCCAAGCGGGGAGTCCGTCACGCTCTCGTTCTCGACTACCGCGGAGAGCCAGTGTGCTGAGGATCCTA 1600
I H D L R A V G A T F A A Q A G A T T K E L M A R L G H T T P R M -

1601 GCGGATGAAGTACCAGATGGCGTCTGAGGCGCGGACGAGGCTATCGCTGAGGCGATGTCCAAGCTGGCCAAAGACCTCTGAAACGCAAAAAGCGCGCGT
Int CCGCTACTTCTATGGTCTACCGCAGACTCCGGGCGGTGCTCGATAGCGACTCCGCTACAGGTTCCGACCGGTTCTGGAGGACTTTCTGTTTTTCCGGGGGA 1700
A M K Y Q M A S E A R D E A I A E A M S K L A K T S -
Int stop

1701 CCCAAGGACACTGAGTCTTAAAGAGGGGGTTTCTTCTCAGTACGCGAAGAACCACGCTTGGCGCGAGCGCGCAGCACCGCGCTCTGTGCGGAGACCTG
GGGTTCTGTGACTCAGGATTCTCTCCCCCAAGAACAGTCAATGCGCTTCTTGGTGGGACCGGGCGCTCGCGGTCTGCGCGGAGACACGCTCTGGAC 1800

PvuII
I

1801 GGCACAGCCCCCGCGCGCCAGGAGCATTGCGGTTCCCGCCAGCTGAGTTCTGTTGTGCGCGCGCTATGTAGAGCTGGTCTGTAGGTCCGATCTCCA
CCGTGGTCCGGGCGGGCGGGTCTCTGTAACGGCAAGGGCGGTGCACTCAAGACAACACGGCGGATACATCTCGACCAGCAACATCCAGGCTAGAGGT 1900

1901 GCGGACTTTCCGGCGACGCTGAGGATGTGATCAGAGCCTCCGGGACCGCGGTTGCGGTCAAACCTGACCATCCGACAGCGGACGCGGTGGTGTCTTC
CCGCTGAAAGGGCGGCTGCGACTCTTACAGCTAGTGTCTCGGAGGCGCTGGCGGCCAAGCGCAGTTTGGACTGGTAGGCTGTGCGCTGCGGCACCAAAAG 2000

SalI
I

2001 CTCCAGGGCTCCGGCTTGCCTGAGAATACAGAGCCAGGTCCCGCTCGGCTCCAGCTCCGACGAGCCCGGTGATCTCTTGGTGCAC 2089
GAGGTCCCGAGGCGGAACGGACTCTTATGTCCTGGTTCGAGGGCGACGCGGAGGTTCGAGGCTGCTCGGGCCACTAGCAGAACCAGCTG






Figure 5




Shaded portion = LS DNA

Figure 6




Shaded portion = L5 DNA

Figure 7



Shaded portion = L5 DNA

Figure 8



Shaded portion = L5 DNA

Figure 9

11





Figure 10



Shaded portion = Kan^R cassette

Figure 11





FIGURE 12



Shaded portion = Kan^R cassette

Figure 13





Figure 14

*Figure 15*




Figure 16




Figure 17




Figure 18





Figure 19

Figure 20






Figure 21




Figure 22




Figure 23




Figure 24

Deleted in p. XVIII, 113-2005

1501
1601
1701
1801
1901
2001
2101
2201
2301
2401
2501
2601

1600
1700
1800
1900
2000
2100
2200
2300
2400
2500
2600
2700

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
81

[illegible]

[illegible]




Figure 23




Figure 26





Figure 27

Synthetic Multiple Cloning Site (MCS) + Strand

GAA GGC GCG GCC GCG GTA CCA GAT CTT TAA ATC TAG ATA TCC ATG GAT
CCA GCT GCA GAA TTC GAA GCT TAT CGA TGT CGA CGT AGT TAA CTA GCG
TAC GAT CGA CTG CCA GGC ATC AAA TAA AAC GAA AGG CTC AGT CGA AAG
ACT GGG CCT TTC GTT TTA TCT GTT GTT TGT CCG GCC ATC ATG GCC GCG
GTG ATC AGC TAG TAC G

Figure 28






Figure 30

pMV206

N
h Begin Kan Cassette
•
•
•
101 GCTAGCCCAACAAAGCGAGCTTGTGTCTCAAAATGTGTGATTTACATTGCGACAAGATAAAAAATATCATCATCAACAAATAAACTGTGTGCTTACATAA
120 GATCGGCTTGTGTGCTGCAACACAGAGTTTACAGACTACAATGTAAAGTGTGTGATTTTATATAGTAGTACTTGTATTTTACAGAGCAATGTATT 130

Kan start codon
101 ACACTAATACAAGGGGTGTTATAGCCATATTCACGGGAACGTGTTGCTGAGGGCGGATTTAAATTCACATGGATGCTGATTTATATGGGTATAA
120 TGTGATTAATGTGTGCGCAATACTGCGTATAAGTGTGCGGTGCGAGAACGAGCTGCGGCGCTAAATTAAGGTTGTACCTACGACTAAATATACCCATATT 130

N
•
•
•
201 ATGGGCTGGGATTAATGTGCGCAATCAGGTGCGACAATGTATGCTTGTATGGGAAGCGGATGCGGCGAGCTTGTGTGTAACATGGCAAGGTAGC
120 TACCGGAGCGCTATTACAGCGCGTTAGTCCAGCGCTGTAGATAGCGAATATACCGTTGCGGCTAGCGGCTGTCAACAAAGACTTGTACCGTTTGTATCG
130

301 GTTGGCAATGATTTACAGATGAGATGGTCAGACTAACTGGGTGACGGAAATTTATGCTTGTGGGACCATCAAGCAATTTATGCGTACTGCTGATGATG
120 CAAGGTTACTACAAATGTCTACTCTACCGAGTGTGATTTGACCGAGCTGCGTTAAATACGGAGAGCGGTGTAGTGTGTAATAAGGCTAGGCTAGGCTACTAC
130

401 CATGCTTACTGACCTGCGATGCGGCGGAAACAGCATTCGAGGTATTAGAGAATATCGTGAATCAGGTGAAATATTTGATGCGGCTGGCAGTGT
120 GTACCAATCAGTGGTGACGCTAGCGGCGCTTTGTGCTAAGGTGCATAATCTTTTATAGGACTAAGTGCAGTTTATACAACTACCGGCGCTACAA
130

501 GGTGCGCGCGTTCGATTCGATTCGCTGTTTAAATGTGCTTTTAAACGGGATCGCTATTTGCTGCTGCTGAGGCGCAATCAGGAATGAATACGGTGT
120 GGACGCGCGCAACGTAAAGGTAAAGGACAACATTAAACAGGAAATTTGCTGAGCGCATAAAGGAGCGAGTCCGCTTGTGTGCTTACTTATTCGCAAC
130

601 GTTGATGCGAGTGATTTTGATGACGAGCTAATGCTGCTGCTTTGAACAGTGTGGAAAGAAATGCATAACTTTTCCATTTGACCGGATTTAGTGG
120 CAACGAGCTTACTAAACTACTGCTGCTGCTAATGCGGAGCACTTTGTCAGACTTTTTCAGCTATTAGAAACGGTAAAGGTGGCTAAGTCCAGC
130

701 TCACTCATGGTGATTTTCTCACTTGATAAGCTTATTTTACGAGCGGAAATTAATAGGTGTGATTTGATTTGGACGAGTGGAAATCGCAGACGGATACCA
120 AGTGAGTACGACTAAAGAGTGAACATTATGGAAATAAACTGCTGCGCTTAAATATCCACATAACTACACGCTGCTCAGCGTTAGCGTGTGCTATGCT
130

801 GGATGTTGCGATGCTATGGAATGCGCTGCTGAGTTTCTGCTTCAATACAGAAACGGCTTTTCAAAAATATGGTATTGATTAATGCTGATATGAATAA
120 GGTAGAACGTTAGGATACCTGACGGAGCGCTCAAAAGAGGAATTAATGCTTTTGGCAAAAGTTTTATACCATAACTATTAGGACTATACTTATT
130

Kan Stop Codon
901 TTGCACTTCACTTTGATGCTGATGAGTTTTCTGATCAGAAATGGTTAAATGGTTTAAACAGTGGCAGAGCATTCAGCTGACTGACGGGAGCGCGCT
120 AACGTCAAAGTAAACTACGAGCTACTCAAAAGATTAGTCTTAAACCAATTAAACCAATGTGACGCTGCTGTAATGGAGTGAAGTGGCTGCGGCGCA
130

1001 TTTTGAAATAATCGAATTTTGTGAGTTGAAGGATCAGATCAGCATCTTCCGACACCGCAGACCGTTCCGTTGGCAAGCAAAAGTCAAAATCACC
120 AACAACTTATTAGCTTGAAACGACTCAACTTCTAGTCTAGTGGTAGAAGGCGTGTGCGTGTGGCAAGGCACCGTTTCTGTTTCAAGTTTATGTTG
130

1101 AACTGCTCAGCTACAAACAAAGCTCTCATCAGCGTGGCTGCTCACTTTCTGCTGATGATGGGGGATTCAGGCTGGTATGAGTCAGCAACACTT
120 TTGACCAAGTGGATGTGTTTGGAGAGTAGTTGGCAGCGAGGAGTGAAGACCGACTACTACCGCGCTAAGTCCGGACCATAGTCACTGCTGTGTGAA
130

S
p Begin E. rep
•
•
•
1201 CTTACAGAGCGAGGCTCAGTATGCTGAGGCTCAGACCGGTAGAAAGATCAAGGATCTTTTGGATCTTTTTTGTGCGGCTAATCTGCTG
1300 GAAGTGTCTGCTGTGAGTGATCAAGGTGACTCGAGTGTGGGCACTTTTCTAGTTTCTGAGAAAGCTTAGGAAAAAAGACCGGCTATTAGACGAC
1301

1301 GTTGAACAAAAAAACACCGCTACCGAGGCTGTTGTTTGGCGGATCAGAGCTAGCAACTTTTTTGGAAAGTAACTGCTTACGAGAGCGCGAG
1400 GACGTTTTGTTTTTGGTGGCGATGCTGGCCACCAACAAACCGGCTAGTTCTGATGCTGAGAAAAAGGCTTCCATTGACCGAAGTCTGCTGCGCTC
1401

1401 ATACCAATATGCTGCTTTAGTGTAGCGGTAGTTAGGCGACCACTTCAAGAACTGTGTAGCAGCGCTACATACCTGCTGCTAATCTGTTTACGAG
1500 TATGTTTTATGACAGGAGATCAATCGGCATCAATCGGTGTTGAAGTCTTTGAGACATCTGGCGGATGTATGAGCGGAGACGATTAGGACAAATGCTC

Figure 31

16301
16302
16303
16304
16305
16306
16307
16308
16309
16310
16311
16312
16313
16314
16315
16316
16317
16318
16319
16320
16321
16322
16323
16324
16325
16326
16327
16328
16329
16330
16331
16332
16333
16334
16335
16336
16337
16338
16339
16340
16341
16342
16343
16344
16345
16346
16347
16348
16349
16350
16351
16352
16353
16354
16355
16356
16357
16358
16359
16360
16361
16362
16363
16364
16365
16366
16367
16368
16369
16370
16371
16372
16373
16374
16375
16376
16377
16378
16379
16380
16381
16382
16383
16384
16385
16386
16387
16388
16389
16390
16391
16392
16393
16394
16395
16396
16397
16398
16399
16400
16401
16402
16403
16404
16405
16406
16407
16408
16409
16410
16411
16412
16413
16414
16415
16416
16417
16418
16419
16420
16421
16422
16423
16424
16425
16426
16427
16428
16429
16430
16431
16432
16433
16434
16435
16436
16437
16438
16439
16440
16441
16442
16443
16444
16445
16446
16447
16448
16449
16450
16451
16452
16453
16454
16455
16456
16457
16458
16459
16460
16461
16462
16463
16464
16465
16466
16467
16468
16469
16470
16471
16472
16473
16474
16475
16476
16477
16478
16479
16480
16481
16482
16483
16484
16485
16486
16487
16488
16489
16490
16491
16492
16493
16494
16495
16496
16497
16498
16499
16500

[illegible]

HSP61 Cassette

1 TCTAGAGGTGACCACAACGACGGCGCCCGCTTTGATCGGGGACGTCTCGCGCCGACCATTACGGGTCTTGTGTCTGTTGGCGGTGATGGCGGCAACATAC 100
CGATCGCCACTGGTGTCTGTCGCGGGGCGAACTAGCCCGTGCAGACGCGGGCTGGTAAATGCCGAGAACACAGCAACCGCCAGTACCGCGCTTGTATG

101 TCACCGGATCGGAGGGCGGAGGACAAAGTCCAAAGAGGGGATGACCGGTGCGGGCTTCTTCCACTCGGCATAGGGGAGTGTAAAGAATAACGTTGG 200
AGTGGGGTACCGTCCCGGCTCTGTCTCCAGCTTGGTCCCGGTACTGGGCGACGCGCGGAAGAAGGTGAGCGGTATCCGCTCAGGATTCTTATTGCAACC

201 CACTCGGACCGGTGAGTCTTAGTTCGGGACGGTGCAGGCCAGGCGCGTCTGTCGAGCGAGTGGCAGCGAGGACAACTTGAGCGCTCGCTCGCGGCGACTG 300
GTGAGCGGTGCGGACTCAGCATCCAGCGCTCCCACTCCGGTCCGGGACGACGGTCTGGTCACCGTCCGCTCTGTGAACTCGGCAGGCGAGCGCGGTGAC

301 CCGCGCGGACGGTAAGTAGCGGGTTCGCGTCCACCGGTGACCGCGGTTTCATCCCGGATCGGAGGAATCACTTCGCAATGGCGAAGACAATTCGCGCGG 400
GGCGCGCGGTCCCATTCATCGCGCGAAGCGGAGTGGGCGACTGGGGGCGAAAGTAGGGGCTAGGGCTCTTAGTGAAAGCGTTACCGGTTCTGTTAAGCGGTAGG
M A K : : A D P

HSP71 Cassette


1 TCTAGACCGCGACGACAGCGTTAGCATGCTCAGTAAGTTGAGTGCATCAGGCTCAGCTCTGAATTGACAGCACACCGCGCTCGAGGCAAGCTTGACGGG 100
AGATCTGGCGGTGCTGGTCCGAATCGTAGGATCACTCAACTCAGCTACTCGAGTCCGAGACTTAACGTCTGTGTCGGCGGACGTCCGTTCCGAATCGCC

101 GGTGCATCATCATAGTGCAGGAAGAAGCTCTACATATTGAGGAGGATTACCATGGCTCTGCGGTGCGGGATCC 177
CCAGGTGAGTAGTATCAGCTCTTCTTCGAGATGTATAAGTCTCTCTAAGTGGTACCGAGCACGCCAGCGCGTAGG
M A R A V G D P




Figure 32

Construction of HSP60 Cassette
PCR of BCG Genomic DNA
HSP60 Primers
Xba-HSP60 + BamHI-HSP61 - HSP261

HSP70 Primers
Xba HSP70 + BamHI-HSP71 - HSP271



PCF110



Digest w/ BamHI
Ligate w/ BamHI *LacZ* cassette
Transform *E. coli*
Select blue colonies on X-gal Plates

Digest w/ BamHI
Ligate w/ BamHI *LacZ* cassette
Transform *E. coli*
Select blue colonies on X-gal Plates





Figure 34







Figure 35



Digest with NotI
Gel Purify 7523 bp fragment
Ligate/Recircularize



Digest with NotI
Gel Purify 7513 bp fragment
Ligate/Recircularize




Figure 36




Figure 37





Figure 38

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US91/04833

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC IPC(5): C12N 15/00, 1/21, 15/11; A61K 39/00 U.S.C1: 536/27; 435/69.1, 252.3; 424/92		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
U.S.	536/27; 435/172.3, 252.3, 69.1; 424/92	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
Databases: Dialog (Files 5, 155, 350, 351, 399), US PTO Automated Patent System (File US Pat., 1975-1991) search terms: mycobacter ? bacteriophage, integrat ?		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹		
Category *	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
<u>X</u> Y	WO, A, 90/00594 (Bloom et al) 25 January 1990, see pages 37-39, 64-70 and Figs. 6 7 .	1, 2, 8-11 3-7
<u>X</u> Y	WO, A, 88/06626 (Bloom et al) 07 September 1988, see entire document.	1, 2, 8-11 3-7
<u>X</u> Y	Proceedings of the National Academy of Sciences, volume 85, issued September 1988, Snapper et al., "Lyso-geny and transformation in mycobacteria: Stable expression of foreign genes", pages 6987-6991, see the entire document.	1, 2, 8-11 3-7
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>* Special categories of cited documents: ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p> </div> </div>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search		Date of Mailing of this International Search Report
15 October 1991		07 NOV 1991
International Searching Authority		Signature of Authorized Officer
ISA/US		 Nancy Treptow

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)

Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
X Y	Reviews of Infectious Diseases volume 11, Supp. 2, issued March-April 1989, Jacobs, Jr. et al., "Myco bacteriophage Vector systems", pages 5404-5410, see entire document.	1, 2, 8-11 3-7
Y	Molecular and General Genetics, volume 195, issued May 1984, Pierson III et al., "Cloning of the integration and attachment regions of bacteriophage P4", pages 44-51, see the entire document.	7
Y	Molecular and General Genetics, volume 192, issued April 1983, Ljungquist et al., "Properties and Products of the cloned <u>int</u> gene of bacteriophage P2," pages 87-94, see the entire document.	7
Y	Journal of Bacteriology, volume 170, No. 12, issued December 1988, Li et al., "Site-specific Integration and expression of a developmental promoter in <u>Myxococcus Xanthus</u> ," pages 5552-5556, see entire document.	7
Y	Journal of Bacteriology, volume 171, No. 3, issued March 1989, Astumian et al., "Site-specific recombination between cloned <u>attP</u> and <u>attB</u> sites from the <u>Haemophilus influenzae</u> bacteriophage HP 1 propagated in recombination-deficient <u>Escherichia coli</u> ," pages 1747-1750, see the entire document.	7